

Review of the clinical and pathological panoply of systemic mastocytosis

Deepti H. Radia,¹  Anna Green,² Clare Oni¹ and Mufaddal Moonim²

¹Department of Clinical Haematology, Guys and St Thomas' NHS Hospitals and ²Department of Cellular Pathology, Guys and St Thomas' NHS Hospitals, London, UK

Summary

Mastocytosis is a rare disease with varied presentation, myriad symptomatology and variable prognosis. Most patients present with cutaneous disease and mediator-related symptomatology with a small subset having systemic disease (systemic mastocytosis, SM). A subset of the latter develops synchronous or metachronous haematologic neoplasms (SM-AHN), most commonly chronic myelomonocytic leukaemia (CMML). Advanced systemic mastocytosis (ASM) is seen in a relatively small number of patients and is usually associated with organ dysfunction, and may present with hepatosplenomegaly, lymphadenopathy and ascites with progression to leukaemic transformation (mast cell leukaemia/acute myeloid leukaemia) occurring in a few patients. This paper discusses the clinical and pathologic features of the entire spectrum of SM in adults.

Keywords: mast cells, systemic mastocytosis, myeloproliferative neoplasm.

Background

Systemic mastocytosis (SM) is a haematological neoplasm with complex diagnostic (Table I) and clinical (Table II) classifications that provide an indication of prognosis and guide treatment (Valent et al., 2001; Swerdlow et al., 2017; Valent et al., 2007; Valent et al., 2017). The spectrum of SM encompasses a heterogeneous population of patients with indolent to aggressive disease and a variable degree of symptom burden (Scherber & Borate 2018) (Fig1). The majority of adult patients (>80%) present with mastocytosis in the skin (MIS), most commonly, maculopapular cutaneous mastocytosis (MPCM), requiring further evaluation to determine if the patient has systemic disease. Utilising the WHO criteria

(Valent et al., 2001; Arber et al., 2016; Valent et al., 2017), patients with a baseline tryptase of >20 µg/l (Schwarz & Irani, 2000) and clinical findings (mediator-related symptoms, musculoskeletal symptoms, constitutional symptoms, organ dysfunction or organomegaly) proceed to biopsy of an extra-cutaneous organ. The bone marrow (BM) biopsy is standard in all patients with a suspected diagnosis of SM. A biopsy of additional organs, such as the liver, gastrointestinal tract, lymph node or body cavity fluid is recommended in those few patients where it is unclear whether the patient has local organ damage and may therefore have a diagnosis of advanced SM. A small minority of patients, who present with severe or life-threatening anaphylaxis following a bee or wasp (Hymenoptera venom) sting or idiopathic anaphylaxis, may also warrant investigation to exclude SM; this subset of patients may have lower baseline tryptase levels (<20 µg/l) and no evidence of cutaneous manifestations (Gulen et al., 2014; Zanotti et al., 2015).

At Guys and St Thomas's Hospitals Foundation Trust, we established a UK centre of excellence in mastocytosis in 2006, in association with the European Competence Network of Mastocytosis (ECNM). We have reviewed more than 500 patients in the mastocytosis service with approximately 200 patients being seen directly within the haematology service. In this review, we will discuss real patient cases to illustrate clinical and histopathological variations in the diagnosis and management of this uncommon disease. Please note patient cases used here span a decade and some pre-2016 terminology for cutaneous mastocytosis will have been applied at the time of diagnosis for some patients.

Given the variable symptomatology, specific tests that are used to identify and manage SM in our routine practice include the evaluation of serum tryptase, liver function tests (LFT), renal function tests (RFT), vitamin D and calcium levels, BM examination (aspirate and trephine), biopsies from other organ systems, molecular analysis (*KIT*, extended myeloid gene panel analysis) and dual-energy X-ray absorptiometry (DEXA) scans. Whole body imaging using MRI or CT scans in patients with SM are useful to assess organomegaly and the presence of lymphadenopathy. In the context of biopsy evaluation, it is pivotal to have access to a facility

Correspondence: Deepti H. Radia, Department of Clinical Haematology, Guy's Hospital, London, SE1 7EH, UK.
E-mail: deepti.radia@gstt.nhs.uk

Table I. Diagnostic criteria for SM (adapted from Swerdlow et al., 2017).

Diagnosis requires:
One major criterion + at least one minor criterion
Or
At least three minor criteria
Major criterion
Multifocal dense infiltrates of mast cells (≥ 15 mast cells in an aggregate) in bone marrow and/or extracutaneous organs
Minor criterion
$>25\%$ of the mast cells in the infiltrate are spindle-shaped or have atypical morphology, or $>25\%$ of all mast cells in bone marrow aspirate smears are immature or atypical
Detection of activating point mutation at codon 816 of <i>KIT</i> in the bone marrow, blood or another extracutaneous organ
Mast cells in bone marrow, blood or another extracutaneous organ express CD25, with or without CD2
Serum total tryptase is persistently >20 ng/ml, unless there is an associated myeloid neoplasm

where appropriate immunohistochemistry can be performed to detect clonal mast cells.

The normal and neoplastic mast cell

Normal mast cells are round, mononuclear cells, with abundant cytoplasm, which contain metachromatic granules in their cytoplasm (Fig 2A). In histological sections, they appear as round cells with abundant pink granulated cytoplasm and small round nuclei (Fig 2B). In healthy patients, a small number of normal mast cells are identified within the BM in a peri-sinusoidal (Fig 2B) or peri-trabecular distribution. They express mast cell tryptase (MCT) and CD117.

Neoplastic mast cells may demonstrate a range of morphologies, from the typical spindle-shaped mast cell with hypogranular cytoplasm (Fig 2C, 2), to those resembling metachromatic blasts. Less frequently, such as in **Case A**, the typical spindle-shaped neoplastic mast cell may show hypergranulation. Bi- or multilobated nuclei and/or blast-like features have been associated with a poorer prognosis (Sperr et al., 2001). Immunophenotypically, in addition to MCT and CD117, neoplastic mast cells co-express CD25 (almost 100%) and CD2 (~60%), although the sensitivity and specificity of assessing CD25 expression alone is comparable to the use of CD2 and CD25 (Jordan et al., 2001; Pardanani et al., 2004; Morgado et al., 2012). In addition, using immunohistochemical staining (rather than flow cytometry), CD25 staining is stronger and easier to assess in our experience. CD2 expression on mast cells is more frequently seen when using flow cytometry. Expression of CD30 and/or CD123 may help in identifying patients who may progress in the future, but data on this association is not particularly robust (Sotlar et al., 2011; Valent et al., 2011; Moonim et al., 2012; Morgado et al., 2013; Doyle et al., 2014; Blatt et al., 2015; Pardanani et al., 2016).

Indolent systemic mastocytosis (ISM)

Case A

A 45-year-old gentleman presented to dermatology with a seven-year history of rashes and facial flushing. He had no history of gastrointestinal symptoms, allergies or anaphylaxis. Clinically, the rash was in keeping with urticaria pigmentosa (now referred to as MPCM) and a skin biopsy confirmed the presence of a dermal mast cell infiltrate (Fig 3A). His tryptase level at diagnosis was raised (136 $\mu\text{g/l}$) with a normal full blood count (Table III). His BM aspirate showed 9% spindle-shaped mast cells. A BM biopsy showed 20% mast cells, which were unusually heavily granulated for mast cells (Fig 3C) and a diagnosis of ISM was made. He was managed with Fexofenadine 180 mg as required and Alendronate for his osteoporosis, and has remained stable during his seven-year follow up with no evidence of disease progression.

Case B (Table IV results summary)

A 44-year-old male patient was referred with a history of four anaphylactic reactions over a period of five weeks. No triggers were identified on review by an Immunologist. He had been diagnosed with UP (now referred to as MPCM) nine years previously and received PUVA treatment. He had mild pruritus and proven allergies to latex, watermelon and shellfish. He suffered from loose bowel motions five to six times a day. No organomegaly was identified. A diagnostic BM aspirate carried out in 2009 showed 4–5% mast cells. The BM biopsy confirmed a diagnosis of SM (disease bulk 5–10%) (Fig 3). Molecular analysis [(allele-specific-polymerase chain reaction (AS-PCR)] did not identify the presence of a *KIT* mutation. He was diagnosed with ISM and treatment was started with sodium cromoglycate 200 mg TDS (three times a day), hydroxyzine hydrochloride 25 mg OD (once daily) and Adcal D3 for his osteoporosis. He carried an Epipen[®] in view of his current presentation. Primary Care Trust (PCT) funding allowed treatment with Imatinib 400 mg OD, which he has remained on for 10 years because of his severe anaphylaxis. No further anaphylactic episodes occurred and he has since stopped his antihistamine and sodium cromoglycate. His tryptase level reduced from 31.1 to 11 $\mu\text{g/l}$ after a year of treatment with Imatinib (Novartis pharmaceuticals).

The pathology of indolent systemic mastocytosis. The diagnosis, in most cases, is made on a BM biopsy. Careful examination of the aspirate often highlights scattered atypical mast cells; but in practicality this can be a time-consuming exercise. A discordance between mast cell burden on BM aspirate and trephine is well-recognised, with the mast cell burden being higher on the BM trephine—the mast cell disease burden is defined by the BM trephine. Diagnosis is made in accordance with the World Health Organisation

Table II. Diagnostic criteria for the variants of SM. Adapted from Swerdlow et al, 2017.

Indolent systemic mastocytosis (ISM)
Meets the general criteria for systemic mastocytosis (Table IIb)
No C findings (see below)
No evidence of an associated haematological neoplasm
Low mast cell burden
Skin lesions invariably present
Smouldering systemic mastocytosis
Meets the general criteria for SM (Table IIb)
≥2 B findings; no C findings (see below)
No evidence of an associated haematological neoplasm
High mast cell burden
Does not meet the criteria for mast cell leukaemia
Systemic mastocytosis with an associated haematological neoplasm (SM-AHN)
Meets the general criteria for SM (Table IIb)
Meets the criteria for an associated haematological neoplasm
Aggressive SM
Meets the general criteria for SM (Table IIb)
≥1 C finding (see below)
Does not meet the criteria for mast cell leukaemia
Skin lesions usually absent
Mast cell leukaemia
Meets the general criteria for SM (Table IIb)
Bone marrow biopsy shows diffuse infiltration by atypical, immature mast cells
Bone marrow aspirate smears show ≥20% mast cells
Leukaemic (classic) mast cell leukaemia: mast cells account for ≥10% of the peripheral white blood cell count
Aleukaemic variant: mast cells account for <10% of the peripheral white blood cell count
B findings
High mast cell burden (on bone marrow trephine biopsy): >30% infiltration of cellularity by mast cells and serum tryptase >200 ng/ml
Signs of dysplasia or myeloproliferation in non-mast cell lineage(s), but criteria are not met for definitive diagnosis of an associated haematological neoplasm, with normal or only slightly abnormal blood counts
Hepatomegaly without impairment of liver function, palpable splenomegaly without hypersplenism and/or lymphadenopathy on palpation or imaging
C findings
Bone marrow dysfunction caused by neoplastic mast cell infiltration ≥1 cytopenia (neutrophils count <1.0 × 10 ⁹ /l, Hb < 100 g/l, platelet count <100 × 10 ⁹ /l)
Palpable hepatomegaly with impairment of liver function, ascites and/or portal hypertension
Skeletal involvement, large osteolytic lesions ± pathological fractures
Palpable splenomegaly with hypersplenism
Malabsorption with weight loss due to gastrointestinal mast cell infiltrates

Clinical Burden of Mastocytosis




Fig 1. Patient-reported symptoms attributed to their systemic mastocytosis according to NIHHR (2016). Note: Intensity of symptoms relative to frequency reported.

Patient-reported symptoms attributed to their SM according to NIHHR (2016). Note: Intensity of symptoms relative to frequency reported.




Fig 2. Normal and neoplastic mast cells. (A) Metachromatic granules (Giemsa 40 \times). (B) Perivascular mast cells (arrows) (H&E 10 \times). (C) Neoplastic mast cell (bone marrow aspirate) (Giemsa 100 \times). (D) Spindle-shaped neoplastic mast cells (H&E 20 \times).

(WHO) major and minor criteria (Swerdlow et al., 2017) (Table 1). The major criterion requires the identification of aggregates of at least 15 mast cells; these aggregates may be distributed as interstitial nodules, as paratrabeular or perivascular aggregates, or may be singly dispersed within the interstitium. In some cases, all patterns may be seen, whereas others may only show interstitial nodules. The distribution within the marrow does not correlate with disease severity. The most frequent cell morphology seen is that of spindle-shaped mast cells, surrounded by a rim of lymphocytes and eosinophils (Figs 3B and 4). Immunophenotypically, CD117 is co-expressed with either CD25 and/or CD2 (Fig 3D, 3 and 3). Well-differentiated systemic mastocytosis (WDSM) should be considered if the mast cells are predominantly round and hypergranulated and show low/negative CD25 expression, as well as lacking *KIT* D816V (Akin, 2004; Alvarez-Twose et al., 2016). These are often responsive to imatinib therapy (Alvarez-Twose et al., 2012).

Practice points: ISM. Allele-specific nested PCR should be used to detect *KIT* D816V in preference to next generation sequencing (NGS), owing to higher sensitivity (sensitivity of allele-specific PCR [between 1–0.01%]; digital droplet PCR [0.005%]).

If a *KIT* D816 mutation is not identified, *KIT* gene sequencing should be considered to assess for any other *KIT* mutation present (exon 17 (D815K, D816F, D816H, D816Y, D820G, E839K), exon 11 (V560G) and exon 10 (K509I, F522C, V530I, A533D)).

Screening for FIP1L1-PDGFR α should be considered if eosinophilia is present.

Anaphylaxis may occur even if BM disease bulk is low.

In the absence of a *KIT* D816V mutation (<10%), Imatinib should be considered as a therapeutic option in symptomatic patients and may lead to complete response in up to 40% patients (Valent et al., 2014; Alvarez-Twose et al., 2016). Patients with mutations in the extracellular/transmembrane domain of *KIT* respond better to imatinib than patients with wild type *KIT*.

Smouldering systemic mastocytosis (SSM)

Case C (Table V results summary)

A 42-year-old male patient was referred to haematology via dermatology in 2010. He was diagnosed with cutaneous mastocytosis at the age of 16 years. He was minimally symptomatic with intermittent pruritus not requiring any medication. In 2008, he sought a dermatology opinion as his rash had significantly worsened, covering most of his body and causing cosmetic concerns, as well as daily pruritus, despite regular loratadine. Dermatology assessment confirmed telangiectasia macularis eruptiva perstans (TMEP) on skin biopsy. He was treated with an attenuated course of PUVA, which was stopped as he did not gain any significant benefit. He had no gastrointestinal symptoms, allergies and had never had any anaphylactic reactions. A diagnostic BM




Fig 3. Indolent systemic mastocytosis (Case A). (A) Skin showing dermal spindle-shaped mast cells (H&E 40 \times). (B) Diagnostic nodular lesion of mastocytosis in bone marrow (H&E 2 \times). (C) Granulated spindle-shaped neoplastic mast cells (Giemsa 40 \times). (D) Mast cell tryptase. (E) CD117. (F) CD2. (G) CD25.

Table III. Results of investigations and clinical findings taken at diagnosis.

Full blood count		
Haemoglobin, g/l	146	
WCC, $\times 10^9/l$	7.1	
Neutrophil count, $\times 10^9/l$	4.8	
Eosinophil, $\times 10^9/l$	0.1	
Platelets, $\times 10^9/l$	225	
Liver and bone profile		
Bilirubin $\mu\text{mol/l}$	9	
ALT IU/L	28	
ALP IU/L	55	
Albumin g/l	50	
Corr. Ca mmol/l	2.36	
Specialist tests		
BMT disease bulk	20%	
Serum tryptase ($\mu\text{g/l}$)	136	
<i>KIT</i>	D816V positive	
Clinical features		
Splenomegaly	None	
Hepatomegaly	None	
DEXA scan	Osteoporosis	

Table IV. Results of investigations and clinical findings taken at diagnosis.

Full blood count		
Haemoglobin, g/l	155	
WCC, $\times 10^9/l$	8.7	
Neutrophil count, $\times 10^9/l$	5.7	
Eosinophil, $\times 10^9/l$	0.3	
Platelets, $\times 10^9/l$	485	
Liver and bone profile		
Bilirubin $\mu\text{mol/l}$	3	
ALT IU/l	45	
ALP IU/l	76	
Albumin g/l	52	
Corr. Ca mmol/l	2.38	
Specialist tests		
BMT disease bulk	5–10%	
Serum tryptase ($\mu\text{g/l}$)	31.2	
<i>KIT</i>	Negative	
Clinical features		
Splenomegaly	None	
Hepatomegaly	None	
DEXA scan	Osteoporosis	

Table V. Results of investigations and clinical findings taken at diagnosis.

Full blood count	
Haemoglobin, g/l	14.6
WCC, $\times 10^9/l$	14.6
Neutrophil count, $\times 10^9/l$	10.5
Eosinophil, $\times 10^9/l$	1.0
Platelets, $\times 10^9/l$	221
Liver and bone profile	
Bilirubin $\mu\text{mol/l}$	6
ALT IU/l	29
ALP IU/l	266
Albumin g/l	51
Corr. Ca mmol/l	2.38
Specialist tests	
BMT disease bulk	30%
Serum tryptase ($\mu\text{g/l}$)	427
<i>KIT</i>	D816V positive
Clinical features	
Splenomegaly	Splenic tip palpable 2 cm below costal margin
DEXA scan	Normal

biopsy was carried out in March 2011 and he was started on fexofenadine 180 mg OD. He was monitored and had two acute admissions for anaphylaxis following a single dose of aspirin (2014) and gastroduodenal bleeds (2015), requiring blood product support and endoscopic laser treatment. These complications of his SM were treated and he is on regular anti-H1 and anti-H2 blockade. Re-evaluation of his disease status in 2015 and 2016 showed an increase in BM disease bulk over time, with accompanying increase in organomegaly evaluated by a CT Scan with increasing abdominal lymphadenopathy and a rising alkaline phosphatase, which peaked at 540 IU/l. Midostaurin was commenced on a compassionate use programme in November 2016 on which he remains well, asymptomatic and with clinical resolution of his organomegaly and TMEP rash.

Pathology. The initial bone marrow transplant (BMT) from March 2011 (Fig 5A–E) shows 30% involvement by SM, in the form of areas of paratrabecular fibrosis, within

which are infiltrates of epithelioid and spindle-shaped mast cells, as well as a dispersed interstitial infiltrate. The mast cells show aberrant expression of CD2 and CD25. The subsequent BM biopsy from July 2016 (Fig 5F, G) shows a slight increase in SM disease bulk to approximately 40%. The mast cells show aberrant expression of CD25, CD2 and CD123, but not CD30. A retroperitoneal lymph node biopsy (Fig 5H–J) performed in July 2016 showed expansion of interfollicular areas by pale, plump, round-to-oval mast cells aberrantly expressing CD25 with heterogeneous expression of CD2, but no expression of CD123 or CD30. A subsequent BM biopsy in October 2016 (Fig 5K–M) showed 50% involvement by SM, in the form of sheets and nodules of round and spindle mast cells expressing CD25 and CD2. The mast cells show focal expression of CD30 and focal expression of CD123.

NB: All his aspirates had <20% mast cells at each biopsy.

This is a typical case of SSM, with high mast cell disease burden (BM disease bulk >30% and serum tryptase >200 ng/ml), splenomegaly and lymphadenopathy.

It has been shown that higher frequency of multilineage involvement of *KIT* D816V in patients with ISM is an independent factor for progression and this may be the aetiology for progression from ISM or SSM for a small number of patients (Escribano et al., 2009).

Case D

A 67-year-old lady presented to dermatology with an 18-month history of erythematous macules affecting her arms, chest and thighs. A skin biopsy confirmed urticaria pigmentosa. She was asymptomatic at the time with no bony pain, gastrointestinal symptoms or mediator symptoms. She had a past history of anaphylactic reaction to alcuronium. There was no lymphadenopathy or hepatosplenomegaly clinically and her baseline results are shown in Table VI. She declined a BM biopsy. She was kept under review and remained asymptomatic. Twelve months from diagnosis the patient developed dyspepsia and flushing symptoms and a BM biopsy showed mast cell infiltrates (45% disease bulk) and evidence of fibrosis. Her peripheral blood film was normal and her BM aspirate showed 10–12% spindle-shaped mast




Fig 4. Low-disease bulk in systemic mastocytosis (SM) (case B). (A) Bone marrow biopsy showing 5–7% involvement by SM. H&E, (B) CD117.




Fig 5. Smouldering systemic mastocytosis (Case C). Bone marrow from 2011, (A). H&E 20 \times . (B) Paratrabecular reticulin fibrosis. (C) CD117 showing 30% disease bulk. (D) CD25. (E) CD2. Bone marrow from July 2016 showing increased disease bulk – 40%. (F) MCT. (G) There is now co-expression of CD123. Retroperitoneal lymph node biopsy from 2016 showing lymph node involvement by mastocytosis. (H). H&E. (I) CD117. (J) CD25. Bone marrow from Oct 2016 showing further increase in disease bulk to 50%. (K) H&E. (M) CD117. (L) The mast cells aberrantly express CD30.

cells. The patient was diagnosed with SSM. Ten years after her diagnosis, this patient continues to remain minimally symptomatic, has undergone successful operations for unrelated medical conditions and now has a palpable spleen. In

this case, high serum tryptase levels correlated with high disease bulk, but this was not associated with a more severe disease course. This case reflects the overlap in spectrum between ISM and SSM.

Table VI. Results of investigations and clinical findings taken at diagnosis.

October 2007	
Full blood count	
Haemoglobin, g/l	129
WCC, $\times 10^9/l$	6.6
Neutrophils $\times 10^9/l$	3.0
Eosinophils, $\times 10^9/l$	0.3
Platelets, $\times 10^9/l$	297
Liver and bone profile	
Bilirubin $\mu\text{mol/l}$	12
ALT IU/l	17
ALP IU/l	156
Albumin g/l	45
Corr. Ca mmol/l	2.22
Specialist tests	
BMT disease bulk	45%
Serum tryptase ($\mu\text{g/l}$)	174
<i>KIT</i>	D816V positive
Clinical features	
Splenomegaly	13 cm on ultrasound
Hepatomegaly	None
DEXA scan	Osteopenia

Pathology (October 2007). The BM shows 45% involvement by paratrabeular and interstitial infiltrates of spindle-shaped mast cells (Fig 6). There is evidence of interstitial fibrosis, with pronounced paratrabeular fibrosis in the areas of mast cell involvement. The mast cells express CD25 and CD2.

Practice points: SSM. Distinguishing SSM from ISM at one end of the spectrum and ASM or mast cell leukaemia (MCL) at the other end needs careful clinical and histological assessment.

Patients with SSM may be asymptomatic or minimally symptomatic, despite the high mast cell burden.

Systemic mastocytosis with an associated haematological neoplasm (SM-AHN)

Introduction. SM is a clonal stem cell disease, and the potential for other lineages to develop clonal aberrations is therefore not surprising. The *KIT* D816V mutation has been clearly demonstrated in cell lineages other than mast cells within the BM of patients with SM (Taylor et al., 2004; Frederiksen et al., 2016). The AHN may precede, be concurrent with or follow the presentation and diagnosis of SM. AHNs span the entire spectrum of haematologic malignancies, including MDS/MPN (myelodysplastic syndromes/myeloproliferative neoplasms) overlap syndromes (CMML being the commonest), MDS, AML, and less commonly lymphomas and plasma cell neoplasms.

Case E

A 60-year-old female was evaluated for erythrocytosis and thrombocytosis following a routine GP review. While a history of allergy to crab meat and penicillin was noted, she did




Fig 6. Smouldering systemic mastocytosis (Case D). Bone marrow biopsy showing paratrabeular zones of fibrosis containing mast cells. (A) H&E 2x. (B) CD117. (C) H&E 20x. (D) Reticulin stain showing increased reticulin fibrosis.

Table VII. Results of investigations and clinical findings taken at diagnosis.

	2005
Full blood count	
Haemoglobin, g/l/PCV	157/0.458
WCC, $\times 10^9/l$	11.7
Neutrophil count, $\times 10^9/l$	7.9
Eosinophil, $\times 10^9/l$	0.1
Platelets, $\times 10^9/l$	510
Liver and bone profile	
Bilirubin $\mu\text{mol/l}$	6
ALT IU/l	19
ALP IU/l	70
Albumin g/l	43
Corr. Ca mmol/l	2.30
Specialist tests	
Disease bulk	<2%
Serum tryptase ($\mu\text{g/l}$)	15
<i>KIT</i>	D816V positive
Clinical features	
Splenomegaly	None
Hepatomegaly	None
DEXA scan	Osteopenia

Table VIII. Results of investigations and clinical findings taken at diagnosis and reassessment.

	Year	
	March 2015	May 2016
Full blood count		
Haemoglobin, g/l	124	138
WCC, $\times 10^9$	7.1	8.0
Neutrophil count, $\times 10^9$	2.9	2.7
Monocyte count, $\times 10^9$	1.5	2.6
Eosinophil, $\times 10^9$	0.4	0.6
Platelets, $\times 10^9$	202	258
Liver and bone profile		
Bilirubin	6	8
ALT	34	14
ALP	241	239
Albumin	45	37
Corr. Calcium	2.21	2.33
Specialist tests		
BMT disease bulk		5–10%
Serum tryptase ($\mu\text{g/l}$)	100	125
<i>KIT</i>		D816V positive
Myeloid gene panel		TET2 mutation allele burden 29%
Clinical features		
Splenomegaly	15.4 cm (ultrasound)	15.5 (on PET)
Hepatomegaly	None	None
DEXA result	Osteopenia	

not have any gastrointestinal or mediator-associated symptoms. Examination showed no rash and no palpable lymphadenopathy or hepatosplenomegaly. Her blood results are shown in Table VII. She was found to be negative for the *JAK2* V617F mutation, had normal ferritin and ESR but a raised carboxyhaemoglobin level of 6.3%. She proceeded to have a BM biopsy which confirmed a diagnosis of essential thrombocythaemia, with an incidental finding of a small clonal mast cell population. A diagnosis of SM-AHN (ISM+ET) was made at that time. She was started on aspirin. A baseline DEXA scan confirmed osteoporosis and she was started on Adcal-D3.

In 2012, seven years after her initial presentation, she developed an urticarial rash which was confirmed as urticaria pigmentosa on skin biopsy; she still had minimal symptoms which infrequently required antihistamine therapy.

This case demonstrates the common association between SM and myeloproliferative neoplasms, and in this case the former was an incidental finding in an originally asymptomatic patient.

Pathology. The BM biopsy from March 2006 (Fig 7A–D) is mildly hypercellular, showing relatively reduced erythropoiesis, with increased megakaryocytes forming loose groups and including some with staghorn nuclei. CD117 staining highlights a focal increase in mast cells (<2% of marrow cells), both spindle-shaped and round forms around bone and blood vessels, with the mast cells showing aberrant expression of CD25. The appearances are of SM-AHN, the AHN component being essential thrombocythaemia.

Case F (Table VIII results)

This 68-year-old lady presented with a five-year history of a rash affecting her face, chest and thighs. A skin biopsy confirmed the presence of mast cells. She reported diarrhoea and lethargy. There was no history of bony pain, flushing or atopy. Clinical examination revealed no lymphadenopathy but there was palpable splenomegaly, which was confirmed on ultrasound. The severity of the patient's diarrhoeal symptoms prompted a colonoscopy, and a colonic biopsy confirmed the presence of mast cells. Her serum tryptase was 100 $\mu\text{g/l}$. A diagnosis of ISM was made.

Thirteen months after diagnosis, the patient developed significant ascites, requiring paracentesis. An upper gastrointestinal endoscopy revealed gastric and duodenal varices. This was accompanied by a rise in her serum tryptase to 125 $\mu\text{g/l}$. A BM examination was performed and showed 5–10% mast cells with evidence of CMML-1. Her monocyte count at this time was 2.6×10^9 . The diagnosis was changed to SM and CMML-1, therefore SM-AHN.

The patient required paracentesis and was treated with steroids for the subsequent two months, at which point midostaurin 100 mg BD (twice daily) was started on a compassionate use programme. She had an excellent clinical

	At trial entry (2012)	Reassessment (2012)	Reassessment (2015)
Full blood count			
Haemoglobin, g/l	84	104	113
WCC, $\times 10^9/l$	5.9	5.3	5.0
Neutrophil count, $\times 10^9/l$	3.4	3.5	3.8
Eosinophil, $\times 10^9/l$	0.2	0.1	0.1
Platelets, $\times 10^9/l$	130	195	148
Liver and bone profile			
Bilirubin $\mu\text{mol/l}$	15	7	8
ALT IU/L	15	19	24
ALP IU/L	154	147	75
Albumin g/l	44	44	49
Corr. Ca mmol/l	2.23	2.22	2.30
Specialist tests			
BMT disease bulk	60–70%	30–35%	40%
Serum tryptase ($\mu\text{g/l}$)	271	137	101
<i>KIT</i>	Positive-D816L		
Clinical features			
Splenomegaly	22 cm*	21 cm*	18.6 cm*
Hepatomegaly	2 cm below costal margin (not recorded on CT)	19.5 cm*	19.2 cm*
DEXA scan	Normal	Normal	Normal

*Craniocaudal measurement via CT.

Table IX. Results of investigations and clinical findings.

response with resolution of ascites, lethargy and significant improvement in her rash, with cessation of steroids. There has not however been any radiological improvement on positron emission tomography (PET) scanning, with no reduction in bowel thickening, abdominal lymph node or spleen size, or bony changes.

In August 2017, she progressed clinically with ascites which was refractory to diuretic treatment, requiring weekly paracentesis and progressive anaemia. A liver biopsy confirmed clonal mast cell infiltration. In November 2017, she consented to be enrolled onto a phase 1 clinical trial using Avapritinib.

Pathology. Biopsies from the colon (Fig 8A–C; October 2015) show a subtle but significant mast cell infiltrate (with sparing of the rectum), including aggregates of >15 mast cells. The infiltrating mast cells express CD25. The subsequent BM (Fig 8D–H; May 2016) is markedly hypercellular, with expanded granulopoiesis, variable eosinophilia, and some megakaryocyte atypia. CD14 highlights increased monocytes. Atypical, spindle-shaped mast cells form peri-trabecular aggregates and account for 5–10% of nucleated marrow cells; the mast cells show aberrant expression of CD25, CD2, CD30 and CD123. The appearances are of SM and CMML1, therefore the pathology fits the category of SM-AHN.

The AHN component in this subtype may be diagnosed concurrently with SM (Case E), or in some cases may

precede or follow the diagnosis of SM (Case F). Where the AHN component is more prominent, it is important not to overlook SM.

The diagnosis of an AHN is prognostically significant, since it determines the patient's prognosis (Naumann et al., 2016). Patients with AHN and additional secondary mutations have a poorer prognosis. Patients who harbour *SRSF2*, *ASXL1* and *RUNX1* mutations in addition to *KIT* D816V ('SAR' mutations) have a significantly reduced overall survival rate (Schwab et al., 2013). The development of recent clinical molecular prognostic models (Naumann et al., 2016; Pardanani et al., 2016; Jahwar et al., 2019) have led to the routine use of myeloid gene panels to help prognosticate patients and guide therapy.

GI Involvement in systemic mastocytosis. Histologically, gastrointestinal involvement by SM shows a broad spectrum of changes (Shih et al., 2016); the disease process may be multifocal, and involvement may be subtle, in which case sampling error is always a possibility. When mast cell burden is low, aberrant CD25 expression is helpful in demonstrating a neoplastic population of mast cells. Aggregates of >15 mast cells in the GI tract fall within the definition of SM (major criterion). However, enumeration of more diffuse infiltrates, and their distinction from a normal population of GI mucosal mast cells may be more difficult. Attempts have been made to quantify mucosal mast cells in SM and diarrhoea. Mucosal mast cells can be increased in non-clonal diarrhoeal

Table X. Results of investigations and clinical findings taken at diagnosis.

Full blood count	
Haemoglobin, g/l	113
WCC, $\times 10^9/l$	18.6
Neutrophil count, $\times 10^9/l$	9.9
Eosinophil, $\times 10^9/l$	4.7
Platelets, $\times 10^9/l$	99
Blood film	Leucoerythroblastic with 1% circulating mast cells.
Liver and bone profile	
Bilirubin $\mu\text{mol/l}$	10
ALT IU/l	55
ALP IU/l	683
Albumin g/l	39
Corr. Ca mmol/l	2.36
Specialist tests	
Bone marrow aspirate	23–25% round mast cells present
BMT disease bulk	40%
Cytogenetics	Normal
Serum tryptase ($\mu\text{g/l}$)	514
<i>KIT</i>	D816V positive
Myeloid gene panel	
GATA2 (47% allele burden)	
SRSF2 (54% allele burden)	
ASXL1 (38% allele burden)	
RUNX1 (44% allele burden)	
Clinical features	
Splenomegaly	Palpable 16 cm below costal margin
Hepatomegaly	Palpable 9 cm below costal margin
DEXA scan	ND

ND, not determined.

illnesses (Doyle et al., 2014), with the mean mast cell infiltrate (per high power field) significantly greater in SM, but this is highly variable and overlaps with non-SM pathologies. Intra-mucosal mast cells may be increased in patients with diarrhoea and IBS-type symptoms; however, confirmation of bowel involvement by SM requires documentation of an immunophenotype normally associated with clonal mast cells.

Practice points: SM-AHN. It is very important to carefully assess the full blood count, differential and background haematopoiesis in all cases of SM to investigate the possibility of an AHN, with CMML being the most common associated haematologic neoplasm.

A myeloid gene panel is useful in evaluating prognosis at baseline in SM-AHN.

In patients where there is doubt over the aetiology – whether it is the SM or AHN component which may be

causing the organ damage – then biopsy of the organ should be considered to guide treatment.

The AHN component often determines the patient's prognosis, and therefore recognition and correct classification is critical to enable appropriate management.

Aggressive mastocytosis (ASM)

Case G (Table IX results). A 26-year-old female was referred in 2012 for consideration of single agent midostaurin in the context of a single arm phase 2 clinical trial for patients with ASM (Gotlib et al., 2016). She had a very unusual history in being diagnosed at the age of four with a persistent rash which had been present since birth, along with hepatomegaly. A diagnosis of urticaria pigmentosa was confirmed on skin biopsy and SM on BM biopsy at the time. She had minimal symptoms needing intermittent antihistamines only and was under regular review. On review in 2011, she was more symptomatic and becoming anaemic. A trial of Interferon α was of little benefit with associated side effects. She was monitored and started to need red cell support to manage her anaemia. She was eligible and consented to start treatment with midostaurin 100 mg BD, to which she had a good clinical response with almost complete resolution of her rash over a three-year period (Fig 9), no palpable organomegaly and normalisation of her blood counts. She remains on midostaurin at present.

The initial BM biopsy in March 2012 (Fig 9D) shows 60–70% involvement by SM, in the form of interstitial nodules and juxta-trabecular aggregates of spindle-shaped mast cells, as well as singly dispersed spindle-shaped mast cells. The mast cells showed aberrant expression of CD25, CD30 and CD123, with focal expression of CD2. There is a change in morphology of the clonal mast cells over time with exposure to midostaurin with loss of clonal markers, as shown in Figs 10 and 11.

Practice points: ASM. Case G has been closely monitored, as required, by participating in a clinical trial of a targeted therapeutic agent. Her repeated BM trephines have shown variable disease bulk over time despite clinical response. Disease burden maybe variable and as sites of BM biopsies vary, the disease burden will too.

The immunophenotype of the mast cells has varied following treatment with midostaurin, raising the possibility of eradication of clones harbouring those markers.

The change in shape from spindle to round hypergranulated mast cells is being noticed more frequently in a post-therapy setting.




Fig 7. SM-AHN [SM-ET] (Case E). (A) Bone marrow showing increased numbers of large megakaryocytes, H&E 4 \times . (B) Hyperlobated megakaryocyte typical of ET, H&E 20 \times . Mast cell aggregates, (C) H&E 10 \times , (D) CD117.

All patients with advanced SM should have a baseline myeloid gene panel.

A holistic approach to interpreting clinical, histopathological and molecular features is needed to guide prognosis and treatment.

In many cases, *KIT*-targeted therapies demonstrated an improvement in cutaneous lesions when seen in ASM, and when the disease responds to the drug.

Mast cell leukaemia (MCL)

Introduction. Mast cell leukaemia is defined by the WHO (Valent et al., 2001; Arber et al., 2016; Valent et al., 2017) as an aggressive variant of SM in which mast cells make up at least 20% of nucleated cells on a *bone marrow aspirate* – these are usually immature morphologically. Two variants are recognised – aleukaemic (peripheral blood mast cells <10%) and leukaemic (peripheral blood mast cells >10%). MCL may arise secondary to existing SM, or may arise *de novo*. The *KIT* D816V mutation is seen less frequently in MCL than in other subtypes of SM (Jawhar et al., 2017). MCL may occur with an AHN, although the reported incidence of this varies in the literature from <10% to >70% (Georgin-Lavialle et al., 2013; Jawhar et al., 2017). While MCL is known to have a very aggressive course (life expectancy <1 year), leukaemic presentations with a more indolent course have been described. These patients lack C-findings

(organ involvement with organ dysfunction) and the mast cells have a more mature appearance on morphology. These patients are labelled as having chronic MCL to differentiate them from the more aggressive variant which is known as acute MCL (Valent et al., 2015).

Case H (Table X results)

This 64-year-old male with MCL was referred for consideration of eligibility for a phase 1 clinical trial. He had no significant past medical history of note and had been previously well. He presented with a three-month history of weight loss (>10% baseline body weight), drenching night sweats, anorexia and severe fatigue. He had noticed a change in his bowel habits with loose stools up to twice a day and an increase in reflux. Examination revealed muscle-wasting and significant hepatosplenomegaly, but no peripheral palpable lymphadenopathy. There was no evidence of cutaneous mastocytosis. This is a case of *de novo* aleukaemic mast cell leukaemia. MCL accounts for <1% of all cases of mastocytosis and is known to have a poor prognosis with a median overall survival of less than six months. He consented to participate in the phase 1 BLU 285 trial for patients with ASM.

Pathology. The diagnosis of aleukaemic MCL was made on the basis of only 1% circulating mast cells in peripheral blood (Fig 12A), the bone marrow aspirate showing 25% immature round granulated mast cells (Fig 12B), and the




Fig 8. SM-AHN progressing to ASM (Case F). Colon (2015) showing lamina propria infiltrates of neoplastic mast cells. (A) H&E, (B) CD117, (C) CD25. Bone marrow (2016) shows a hypercellular marrow with increased monocytes (CMML-1). (D, E) H&E, (F) CD14. Mast cell aggregates with fibrosis are clearly seen. (G) H&E, (H) CD117.

bone marrow trephine biopsy showing hypercellularity, with around 40% of nucleated marrow cells being mast cells. The majority of the mast cells are round, arranged either as interstitial aggregates or singly dispersed within the interstitium. They show aberrant expression of CD25, CD30 with patchy weak expression of CD123.

Practise points: mast cell leukaemia. The bone marrow aspirate is pivotal in making a diagnosis of MCL (>20% mast cells).

MCL can be aleukaemic (peripheral blood <10% mast cells) or leukaemic (>10% circulating mast cells).

Acute MCL is defined by immature morphology and an aggressive course. Chronic MCL is defined by mature mast cell morphology and an indolent course.

Mast cells in MCL tend to be immature and lose the spindle shape conventionally seen in ISM.

Morphologic variants include a spectrum from blast-like, round to spindle-shaped cells with varying degrees of granularity. Some forms may have coarse purple granules (Georgin-Lavialle et al., 2013; Valent et al., 2015; Swerdlow et al., 2017).

Patients with MCL have the poorest overall survival (Paradanani et al., 2016) and should be considered for trials using newer targeted agents, high dose chemotherapy and bone marrow transplant if clinically eligible.

Bone marrow pathology summary

The percentage of neoplastic mast cells in the BM (compared to all nucleated marrow cells), has been shown to




Fig 9. Aggressive systemic mastocytosis (Case G). Bone marrow aspirate showing clusters of mast cells. (A) (low power), (B) (high power). The mast cells are an admixture of spindle-shaped and round hypergranular forms (C). Bone marrow biopsies showing variable disease bulk across the years. (D) March 2012, 70%. (E) Sept 2012, 30%. (F) Dec 2015, 40%.

correlate with disease prognosis (Sperr et al., 2001). However, we have found that BM disease bulk is not always indicative of the patient’s current clinical parameters. This highlights the fact that subtyping of SM is not possible on the basis of the BMT findings alone. Given that SM can show a variable distribution in the BM, sampling bias should always be considered if such a discrepancy arises, and the whole clinical picture taken into consideration, especially if there are other known sites of SM involvement.

It can be difficult to make a diagnosis if the neoplastic mast cells show only a paratrabecular distribution, with significant associated fibrosis (Case D).

CD25 and CD2 immunohistochemical staining is routinely used to identify aberrant expression in neoplastic mast cells, and their expression (CD25 and/or CD2) form one of the minor criteria for the diagnosis of SM (Swerdlow et al., 2017).

Although the typical appearance of SM in the BM is that of aggregates of spindle-shaped mast cells, occasionally the




Fig 10. Bone marrow disease bulk over the years for case G, along with expression profile of mast cell markers by immunohistochemistry (green: expressed, red: negative).




Fig 11. Aggressive systemic mastocytosis (Case G). Skin disease (A) Pre-treatment. (B) After treatment with midostaurin.




Fig 12. Mast cell leukaemia (Case H). (A) Circulating mast cell in peripheral blood (Oil 60 \times). (B) Immature round granulated mast cells in bone marrow aspirate (Oil 60 \times). (C) Hypercellular bone marrow biopsy. Nodules (D) and singly scattered round mast cells (E) within the bone marrow (mast cell tryptase IHC).

predominant feature may be singly dispersed mast cells, which are morphologically atypical with aberrant expression of CD25, highlighting the importance of immunohistochemical staining in identifying such a population.

In addition, aberrant expression of CD30 and CD123 have been increasingly recognised throughout the spectrum of SM

pathologies. Both CD30 and CD123 expression has been reported to be associated with more aggressive disease (Sotlar et al., 2011; Valent et al., 2011; Blatt et al., 2015). This prognostic significance is not consistently reported in the literature (Morgado et al., 2013; Doyle et al., 2014) and in our experience, in a cohort of 43 patients (Moonim et al., 2012),

Overview of current treatment options for patients with systemic mastocytosis 2019.




Fig 13. Overview of current treatment options for patients with SM 2019. *Combinations may be needed and doses higher than those recommended in the British National Formulary. **Midostaurin has been approved by the FDA/EMA for patients with advanced SM. It is currently under review by NICE in the UK.

we did not find direct association between CD30 and CD123 expression and aggressive behaviour. However, Moonim et al. (2012) did identify that CD30 expression correlated with higher disease bulk. In our practice, we have seen variability in aberrant antibody expression between sites of disease and over time. **Case C** (SSM) showed expression of CD123 but not CD30 in the first BMT, no expression of either antibody in the lymph node involved by SM, and focal expression of both antibodies in a subsequent biopsy.

Pathology Practice Points

Although neoplastic mast cells are typically hypogranulated, hypergranulation may be seen, and does not exclude a diagnosis of SM.

Paratrabeular aggregates of mast cells can be associated with significant fibrosis, which may distract from the identification of neoplastic mast cells, especially when disease burden is low.

The distribution of neoplastic mast cells within the BM does not correlate with disease severity.

Although BM disease bulk has been shown to correlate with prognosis, this is not absolute, and does not distinguish SM subtypes.

ISM may show a highly variable distribution in the BM, sampling bias should always be considered if there is any discrepancy between BM findings and the full clinical picture.

KIT mutation testing at diagnosis should be carried out by allele-specific nested PCR technique as this is more sensitive than NGS.

NGS-based myeloid gene panels should be carried out in all cases of advanced SM. Additional somatic mutations are seen in approximately 90% of patients with advanced SM and predominantly in those with SM-AHN. The commonly seen mutations are TET2, SRSF2, ASXL1, EZHZ, CBL, RUNX1.

CONCLUSIONS

Systemic mastocytosis is a rare haematological neoplasm with a spectrum of clinical manifestations, often unique to each patient. The diagnostic and classification pathways are complex. They highlight the need for accurate application of diagnostic criteria, setting up of diagnostic pathways, including molecular analysis with review of histology by experienced haematopathologists and management by clinicians with expertise in specialist comprehensive care centres. The current treatment options for patients with SM are

summarised in Fig 13. With novel targeted therapies entering clinical trial arenas, there is now hope for this orphan group of patients who frequently have a poor prognosis or severely impaired quality of life. It is vital we continue to work internationally and collaboratively to improve our understanding of the pathophysiology of this disease, impact of treatments and move towards truly personalised treatment.

Acknowledgements

We would like to acknowledge the following members of our mastocytosis team: Dr C. Grattan, Professor C. N. Harrison,

Dr D. McLornan, Dr N. Curto-Gracia, C. Woodley, Y. Frances, Dr Y. Miki, Dr M Ong, Dr B. Wilkins, Dr J. van der Walt, D. Drasutyte, M. Ciesielska, T. Olaoni and Professor N. C. P. Cross.

Author Contribution

All four authors DHR, AG, CO and MTM were responsible for all aspects of research, article preparation and writing.

Conflicts of Interest

No conflicts of interest exist.

References

- Akin, C. (2004) Well-differentiated systemic mastocytosis*1A new disease variant with mature mast cell phenotype and lack of codon 816 c-Kit mutations. *Journal of Allergy and Clinical Immunology*, **113**, S327.
- Alvarez-Twose, I., González, P., Morgado, J.M., Jara-Acevedo, M., Sánchez-Muñoz, L., Matito, A., Mollejo, M., Orfao, A. & Escribano, L. (2012) Complete response after imatinib mesylate therapy in a patient with well-differentiated systemic mastocytosis. *Journal of Clinical Oncology*, **30**, e126–e129.
- Álvarez-Twose, I., Jara-Acevedo, M., Morgado, J.M., García-Montero, A., Sánchez-Muñoz, L., Teodósio, C., Matito, A., Mayado, A., Caldas, C., Mollejo, M., Orfao, A. & Escribano, L. (2016) Clinical, immunophenotypic, and molecular characteristics of well-differentiated systemic mastocytosis. *Journal of Allergy and Clinical Immunology*, **137**, 168–178.e1.
- Alvarez-Twose, I., Matito, A., Morgado, J.M., Sánchez-Muñoz, L., Jara-Acevedo, M., García-Montero, A., Mayado, A., Caldas, C., Teodósio, C., Muñoz-González, J.I., Mollejo, M., Escribano, L. & Orfao, A. (2016) Imatinib in systemic mastocytosis: a phase IV clinical trial in patients lacking exon 17 KIT mutations and review of the literature. *Oncotarget*, **8**, 68950–68963.
- Arber, D.A., Orazi, A., Hasserjian, R., Thiele, J., Borowitz, M.J., Le Beau, M.M., Bloomfield, C.D., Cazzola, M. & Vardiman J.W. (2016) The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*, **127**, 2391–2405.
- Blatt, K., Cerny-Reiterer, S., Schwaab, J., Sotlar, K., Eisenwort, G., Stefanzi, G., Hoermann, G., Mayerhofer, M., Schneeweiss, M., Knapp, S., Rüllicke, T., Hadzijusufovic, E., Bauer, K., Smiljkovic, D., Willmann, M., Reiter, A., Horny, H.P. & Valent, P. (2015) Identification of the Ki-1 antigen (CD30) as a novel therapeutic target in systemic mastocytosis. *Blood*, **126**, 2832–2841.
- Doyle, L.A., Sepehr, G.J., Hamilton, M.J., Akin, C., Castells, M.C. & Hornick, J.L. (2014) A clinicopathologic study of 24 cases of systemic mastocytosis involving the gastrointestinal tract and assessment of mucosal mast cell density in irritable bowel syndrome and asymptomatic patients. *The American Journal of Surgical Pathology*, **38**, 832–843.
- Escribano, L., Alvarez-Twose, I., Sanchez-Munoz, L., Garcia-Montero, A., Nunez, R., Almeida, J., Jara-Acevedo, M., Teodosio, C., Garcia-Cosio, M., Bellas, C. & Orfoa, A. (2009) Prognosis in adult systemic mastocytosis; a long-term study of the Spanish Network on Mastocytosis in a series of 145 patients. *The Journal of Allergy and Clinical Immunology*, **124**, 514–521.
- Frederiksen, J.K., Shao, L., Bixby, D.L. & Ross, C.W. (2016) Shared clonal cytogenetic abnormalities in aberrant mast cells and leukemic myeloid blasts detected by single nucleotide polymorphism microarray-based whole-genome scanning. *Genes, Chromosomes and Cancer*, **55**, 389–396.
- Georgin-Lavialle, S., Lhermitte, L., Dubreuil, P., Chandresris, M.O., Hermine, O. & Damaj, G. (2013) Mast cell leukaemia. *Blood*, **121**, 1285–1295.
- Gotlib, J., Kluijn-Nelemans, H.C., George, T.I., Akin, C., Sotlar, K., Hermine, O., Awan, F.T., Hexner, E., Mauro, M.J., Sternber, D.W., Villeneuve, M., Labed, A.H., Stanek, E.J., Hartmann, K., Horny, H.P., Valent, P. & Reiter, A. (2016) Efficacy and safety of midostaurin in advanced systemic mastocytosis. *New England Journal of Medicine*, **374**, 2530–2541.
- Gulen, T., Hagglund, H., Dahken, B. & Nilsson, G. (2014) High prevalence of anaphylaxis in patients with systemic mastocytosis - a single-centre experience. *Clinical & Experimental Allergy*, **44**, 121–129.
- Jawhar, M., Schwaab, J., Meggendorfer, M., Naumann, N., Horny, H.P., Sotlar, K., Haferlach, T., Schmitt, K., Fabarius, A., Valent, P., Hofmann, W.K., Cross, N.C.P., Metzgeroth, G. & Reiter, A. (2017) The clinical and molecular diversity of mast cell leukemia with or without associated hematologic neoplasm. *Haematologica*, **102**, 1035–1043.
- Jawhar, M., Schwaab, J., Alvarez-Twose, I., Shoumariyeh, K., Naumann, N., Lubke, J., Perkins, C., Munoz-Gonzalez, J., Meggendorfer, M., Kennedy, V., Metzgeroth, G., Fabarius, A., Pfeifer, D., Sotlar, K., Horny, H.-P., Bubnoff, N., Haferlach, T., Cross, N.C.P., Hpfmann, W.-K., Sperr, W., Garcia-Montero, A., Valent, P., Gotlib, J., Orfoa, A. & Reiter, A. (2019) MARS: mutation-adjusted risk score for advanced systemic mastocytosis. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*. JCO1900640. <https://doi.org/10.1200/JCO.19.00640>. Oral presentation EHA, Amsterdam. 16th June 2019.
- Jordan, J.H., Walchshofer, S., Jurecka, W., Mosberger, I., Sperr, W.R., Wolff, K., Chott, A., Bühring, H.J., Lechner, K., Horny, H.P. & Valent, P. (2001) Immunohistochemical properties of bone marrow mast cells in systemic mastocytosis: evidence for expression of CD2, CD117/Kit, and bcl-xL. *Human Pathology*, **32**, 545–552.
- Moonim, M.T., Kossier, T., van Der Walt, J., Wilkins, B., Harrison, C.N. & Radia, D.H. (2012) CD30/CD123 expression in systemic mastocytosis does not correlate with aggressive disease. *Blood*, **120**, 1746–1746.
- Morgado, J.M., Sánchez-Muñoz, L., Teodósio, C.G., Jara-Acevedo, M., Alvarez-Twose, I., Matito, A., Fernández-Nuñez, E., García-Montero, A. & Orfao, A. (2012) Immunophenotyping in systemic mastocytosis diagnosis: 'CD25 positive' alone is more informative than the 'CD25 and/or CD2' WHO criterion. *Modern Pathology*, **25**, 516–521.
- Morgado, J.M., Perbellini, O., Johnson, R.C., Teodósio, C., Matito, A., Álvarez-Twose, I., Bonadonna, P., Zamò, A., Jara-Acevedo, M., Mayado, A., Garcia-Montero, A., Mollejo, M., George, T.I., Zanotti, R., Orfao, A., Escribano, L. & Sánchez-Muñoz, L. (2013) CD30 expression by bone marrow mast cells from different diagnostic variants of systemic mastocytosis. *Histopathology*, **63**, 780–787.
- Naumann, N., Jawhar, M., Schwaab, J., Metzgeroth, G., Popp, H., Khaled, N., Horny, H.P., Sotlar, K., Valent, P., Haferlach, C., Gohring, G., Schlegelberger, B., Meggendorfer, M., Cross, N.C.P., Hofmann, W.-K., Reiter, A. & Fabarius,

- A. (2016) Incidence and prognostic impact of cytogenetics in combination with molecular aberrations in patients with systemic mastocytosis. *Blood*, **128**, 947.
- Pardani, A., Kimlinger, T., Reeder, T., Li, C.Y. & Tefferi, A. (2004) Bone marrow mast cell immunophenotyping in adults with mast cell disease: a prospective study of 33 patients. *Leukemia Research*, **28**, 777–783.
- Pardani, A., Reichard, K.K., Zblewski, D., Abdelrahman, R.A., Wassie, E.A., Morice Ii, W.G., Brooks, C., Grogg, K.L., Hanson, C.A., Tefferi, A. & Chen, D. (2016) CD123 immunostaining patterns in systemic mastocytosis: differential expression in disease subgroups and potential prognostic value. *Leukemia*, **30**, 914–918.
- Scherber, R.M. & Borate, U. (2018) How we diagnose and treat systemic mastocytosis in adults. *British Journal of Haematology*, **180**, 11–23.
- Schwaab, J., Schnittger, S., Sotlar, K., Walz, C., Fabarius, A., Pfirmann, M., Kohlmann, A., Grossmann, V., Meggendorfer, M., Horny, H.P., Valent, P., Jawhar, M., Teichmann, M., Metzgeroth, G., Erben, P., Ernst, T., Hochhaus, A., Haferlach, T., Hofmann, W.K., Cross, N.C. & Reiter, A. (2013) Comprehensive mutational profiling in advanced systemic mastocytosis. *Blood*, **122**, 2460–2466.
- Schwarz, L.B. & Irani, A.M. (2000) Serum tryptase and laboratory diagnosis of systemic mastocytosis. *Hematology/Oncology Clinics of North America*, **14**, 641–657.
- Shih, A.R., Deshpande, V., Ferry, J.A. & Zukerberg, L. (2016) Clinicopathological characteristics of systemic mastocytosis in the intestine. *Histopathology*, **69**, 1021–1027.
- Sotlar, K., Cerny-Reiterer, S., Petat-Dutter, K., Hessel, H., Berezowska, S., Müllauer, L., Valent, P. & Horny, H.-P. (2011) Aberrant expression of CD30 in neoplastic mast cells in high-grade mastocytosis. *Modern Pathology*, **24**, 585–595.
- Sperr, W.R., Escibano, L., Jordan, J.H., Scherthaner, G.H., Kundi, M., Horny, H.P. & Valent, P. (2001) Morphologic properties of neoplastic mast cells: delineation of stages of maturation and implication for cytological grading of mastocytosis. *Leukemia Research*, **25**, 529–536.
- Swerdlow, S.H., Campo, E., Harris, N.L., Jaffe, E.S., Pileri, S.A., Stine, H., Thiele, J., Arber, D.A., Hasserjian, R.P., Le Beau, M.M., Orazi, A. & Siebert, R. (eds.) (2017) WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. IARC, Lyon.
- Taylor, M.L., Sehgal, D., Raffeld, M., Obiakor, H., Akin, C., Mage, R.G. & Metcalfe, D.D. (2004) Demonstration that mast cells, T cells, and B cells bearing the activating kit mutation D816V occur in clusters within the marrow of patients with mastocytosis. *The Journal of Molecular Diagnostics*, **6**, 335–342.
- Valent, P., Horny, H.P., Escibano, L., Longley, B.J., Li, C.Y., Schwartz, L.B., Marone, G., Nuñez, R., Akin, C., Sotlar, K., Sperr, W.R., Wolff, K., Brunning, R.D., Parwaresch, R.M., Austen, K.F., Lennert, K., Metcalfe, D.D., Vardiman, J.W. & Bennett, J.M. (2001) Diagnostic criteria and classification of mastocytosis: a consensus proposal. *Leukemia Research*, **25**, 603–625.
- Valent, P., Akin, C., Escibano, L., Födinger, M., Hartmann, K., Brockow, K., Castells, M., Sperr, W.R., Klun-Nelemans, H.C., Hamdy, N.A., Lortholary, O., Robyn, J., van Doormaal, J., Sotlar, K., Hauswirth, A.W., Arock, M., Hermine, O., Hellmann, A., Triggiani, M., Niedoszytko, M., Schwartz, L.B., Orfao, A., Horny, H.P. & Metcalfe, D.D. (2007) Standards and standardization in mastocytosis: consensus statements on diagnostics, treatment recommendations and response criteria. *European Journal of Clinical Investigation*, **37**, 435–453.
- Valent, P., Sotlar, K. & Horny, H.P. (2011) Aberrant expression of CD30 in aggressive systemic mastocytosis and mast cell leukemia: a differential diagnosis to consider in aggressive hematopoietic CD30-positive neoplasms. *Leukemia & Lymphoma*, **52**, 740–744.
- Valent, P., Cerny-Reiterer, S., Hoermann, G., Sperr, W.R., Müllauer, L., Mannhalter, C. & Pehamberger, H. (2014) Long-lasting complete response to imatinib in patient with systemic mastocytosis exhibiting wild type KIT. *American Journal of Blood Research*, **4**, 93–100.
- Valent, P., Sotlar, K., Sperr, W.R., Reiter, A., Arock, M. & Horny, H.P. (2015) Chronic mast cell leukemia: a novel leukemia-variant with distinct morphological and clinical features. *Leukemia Research*, **39**, 1–5.
- Valent, P., Akin, C. & Metcalfe, D. (2017) Mastocytosis: 2016 updated WHO classification and novel emerging treatment concepts. *Blood*, **129**, 1420–1427.
- Zanotti, R., Lombardo, C., Passalacqua, G., Caimmi, C., Bonifacio, M., De Giovanna, M., Perbellin, O., Rossini, M., Schena, D., Busa, M., Marcotulli, M.C., Bilo, M.B., Franchini, M., Marchi, G., Livio, S. & Bonadonna, P. (2015) Clonal mast cell disorder in patients with severe Hymenoptera venom allergy and normal serum tryptase. *Journal of Allergy and Clinical Immunology*, **136**, 135–139.